Chemical Components of *Ocimum basilicum* L. and *Ocimum tenuiflorum* L. Stem Essential Oils and Evaluation of Their Antioxidant Activities Using DPPH Method

Ni Putu Ermi Hikmawanti, Hariyanti*, Nurkamalia, Sahera Nurhidayah

Faculty of Pharmacy and Sciences, Universitas Muhammadiyah Prof. DR. HAMKA, Jakarta, Indonesia

ABSTRACT

Ocimum basilicum L. and *Ocimum tenuiflorum* L. are two types of plants from *Ocimum* (Lamiaceae). One of their chemical compounds is essential oil. Essential oil might have an antioxidant activities. The part of the plants which are often wasted and never used are stem. This study aimed to analyze the chemical components of *O. basilicum* L. and *O. tenuiflorum* L. stem essential oils and evaluate their antioxidant activities. Essential oils were obtained by steam-water distillation. Analysis of chemical component of essential oil was performed using Gas Chromatography-Mass Spectroscopy (GC-MS). Antioxidant activities were evaluated with DPPH method and then the IC₅₀ value was determined as 50% inhibition concentration of free radical. The results showed that *O. basilicum* L. stem essential oil had 13 components with the major compounds were methyl eugenol (52.60%), caryophyllene (18.75%), and germacrene-D (9.19%). Whereas, *O. tenuiflorum* L. stem essential oil had 11 components with the major compounds were α -copaene (5.56%), caryophyllene (17.28%), germacrene-D (9.29%) and methyl eugenol (56.72%). IC₅₀ value of *O. basilicum* L. stem essential oil was 17.50 μ g/mL, whereas IC₅₀ value of *O. tenuiflorum* L. stem essential oil was 14.17 μ g/mL. It was concluded that both oils might be good natural antioxidant agents.

ARTICLE HISTORY

Received: June 2019

Revised: August 2019

Accepted: September 2019

*Corresponding author

Email: hariyanti.irawan@gmail.com

Keywords: antioxidant; essential oils; *Ocimum basilicum* L.; *Ocimum tenuiflorum* L.; stem

INTRODUCTION

Four or five species of *Ocimum* spp. grow in Indonesia. Two of the five species are distinct and should be relatively easy to identify. Whereas, the other two (or three) species are more similar in appearance. *O. basilicum* L. can be identified because it has the largest flowers of any basil found in Indonesia. Stems and leaves are glabrous (smooth, without hairs) or only minutely hairy. *O. tenuiflorum* L. is often encountered under the synonym *O. sanctum* L. It has been described as a short-lived perennial and a dwarf shrub (or subshrub). It is much branched (Mead, 2014).

Ocimum tea is efficacious to expel gases, overcome stomach cramps, constipation, diarrhea and vomiting. It is also useful for treating mental fatigue, nervous conditions and hyssop for cough (Marwat et al., 2011). Indonesian people mostly use leaves of *O. basilicum* L. as fresh vegetables. *O. tenuiflorum* L. are also used as a spice in cooking because the aroma can reduce the fishy smell of wet fish. While the stem of this plant is often wasted and not used.

O. basilicum L. plant contains essential oils, triterpene, alkaloids, flavonoids, saponins, coumarin, steroids, glycoside and tannins. The essential oils contain monoterpene hydrocarbons, oxygenated monoterpene, sesquiterpene hydrocarbons and oxygenated

sesquiterpene (Marwat et al., 2011). Whereas, the main chemical components in *O. tenuiflorum* L. are oleanolic acid, ursolic acid, rosmarinic acid, eugenol, carvacol, linalool and caryophyllene (Ravi et al., 2012).

Essential oils are natural plant products that have various biological properties. Essential oils contain volatile compounds (mainly mono- and sesquiterpenoids, benzenoids, phenylpropanoids, etc.) (Baser and Buchbauer, 2010). This oil is produced from certain parts of plant, such as roots, stems, skin, leaves, flowers and seeds (Gunawan and Mulyani 2004; Lutony and Rahmayati, 1994). Essential oils extracted from *Ocimum* plants have been applied to inhibit growth of microorganisms, in perfumery, in food preservation and also in aromatherapy (Pandey et al., 2014). A large number of studies on the antioxidant potential of essential oils are aimed at obtaining natural non-toxic antioxidants. Several compounds, such as eugenol, α -pinene, terpinene, phellandrene, etc., are example of chemical components of essential oils that are thought to be related to the antioxidant activity (Amorati et al., 2013).

Studying volatile constituent components can be done by profiling the oil metabolites using the gas chromatography-mass spectrometry method. In addition, the process of determining secondary metabolites with certain characteristics is related to the response and mechanism of a drug (Saifudin et al., 2011). Thus, this

study aimed to analyze the chemical components of *O. basilicum* L. and *O. tenuiflorum* L. stem essential oils and evaluate their antioxidant activities.

METHODS

Plant Collection and Identification

Samples were collected from the Indonesian Spices and Medicinal Crops Research Institute (ISMCR)/Balai Penelitian Tanaman Rempah dan Obat (BALITTRO), Bogor, West Java. All samples then were identified in Herbarium Bogoriense, Biology Research Center, Indonesian Institute of Sciences/Lembaga Ilmu Pengetahuan Indonesia (LIPI), Cibinong.

Extraction of Essential Oils

The extraction process was carried out in the BALITTRO, Bogor, West Java. 12 kg of fresh *O. basilicum* L. stem and 20 kg of fresh *O. tenuiflorum* L. stem were extracted separately using the steam-water distillation for 6 hours. Essential oils were obtained then added with anhydrate Na₂SO₄ to obtain pure essential oils. Essential oils were then analyzed for their characteristics including organoleptic parameters, yield percentages, specific gravity, refractive index and solubility in absolute alcohol.

Analysis of Essential Oils

Analysis of essential oils was carried out in the BALITTRO, Bogor, West Java. 10 µL of the *Ocimum* essential oil were mixed with 1 mL of GC grade n-hexane. One µL of the mixture was analyzed with the Gas Chromatography with Auto Sampler (Agilent Technologies 7890), 5975 Mass Selective Detector and Chemstation data system. Separation was carried out using the capillary column (Innowax HP) with length of 30 m x 0.25 mm and film thickness of 0.25 µm. The carrier gas was helium gas at a constant flow rate of 0.6 mL/min. The injector temperature was set at 250 °C. The initial temperature of the oven was set at 60 °C then hold for 0 min, rising at 2 °C/min to 150 °C then hold for 1 min and finally rising at 20 °C/min to 210 °C then hold for 10 min. According to Herebian *et al.* (2009), the main compounds in the GC-MS chromatogram have criteria for percent area >5% of essential oils. The similarity of mass spectra of sample with a library was determined at a qualifier value of at least 80%. The components was identified based on compatibility with an authentic mass spectrum in Wiley electronic library.

DPPH Radical Scavenging Activity

The free radical scavenging activity assay using the 1,1-diphenyl-2-picryl-hydrazyl (DPPH) reagent was determined according to Brand-Williams *et al.* (1995). 0.3 mL of 100 µg/mL of essential oil was reacted with 0.5 mL of DPPH 0.1 mM solution in ethanol and 1.5 mL

of absolute ethanol. The solution was incubated in the dark at room temperature for 30 min. The decolorizing (from deep violet to light yellow) process was read at 516 nm using UV-VIS Spectrophotometer (Shimadzu) and compared with a blank control. For comparison, activities of Vitamin E standard was evaluated.

In the analytical procedure, the measurements were performed at four points of concentration, including control. The measurement of the DPPH radical scavenging activity of sample solution was repeated three times at each concentration. The results were expressed as percentage of decrease of DPPH radical. DPPH inhibition percentage is calculated using the following formula:

$$\text{Inhibition ratio (\%)} = \frac{(\text{absorbance of control} - \text{absorbance of sample})}{\text{absorbance of control}} \times 100 \%$$

The sample concentrations (x) were plotted against the inhibition ratios (y) and then the regression line equation ($y = bx \pm a$) was obtained. The calculation of 50% of the inhibition concentration of DPPH radical is obtained by entering 50 on y in the regression line equation, so x is the IC₅₀ value.

RESULTS AND DISCUSSION

Extraction yields of *O. basilicum* L. and *O. tenuiflorum* L. stem essential oils are shown in Table 1. Oil contents (% v/w) in fresh stems of *O. basilicum* L. and *O. tenuiflorum* L. were 0.183% and 0.200%, respectively. Characteristics of essential oils are shown in Table 2. Chromatograms of *O. basilicum* L. and *O. tenuiflorum* L. stem essential oils are shown in Figure 1 and Figure 2. *O. basilicum* L. stem essential oil had 13 components. The major chemical constituents (>5%) from this oil were methyl eugenol (52.60%), caryophyllene (18.75%) and germacrene-D (9.19%). Whereas, *O. tenuiflorum* L. stem essential oil had 11 components with the major compounds were α-copaene (5.56%), caryophyllene (17.28%), germacrene-D (9.29%) and methyl eugenol (56.72%). Chemical constituents of the essential oils are shown in Table 3. IC₅₀ values of DPPH radical from *O. basilicum* L. and *O. tenuiflorum* L. stem essential oils were 17.50 µg/mL and 14.17 µg/mL, respectively. Whereas, the IC₅₀ value of Vitamin E was obtained at 30.89 µg/mL.

In term of composition, essential oil of the stem is different from essential oil of aerial parts. Essential oils from *O. basilicum* L. aerial parts from Cianjur, West Java were reported to contain 4.88% methyl eugenol. On the other hand, the essential oil of *O. tenuiflorum* L. has a content of methyl eugenol of 8.69% (Sulianti, 2008). *O. basilicum* essential oil from China, Croatia, Israel, Republic of Guinea, Nigeria, Egypt, Pakistan and Malaysia was reported having a major component

Table 1. Extraction yield of *O. basilicum* L. and *O. tenuiflorum* L. stem essential oils

Sample	Weight of fresh stem (Kg)	Volume of essential oil (mL)
<i>O. basilicum</i> L	12	22
<i>O. tenuiflorum</i> L	10	20

Table 2. Characteristics of *O. basilicum* L. and *O. tenuiflorum* L. stem essential oils

Parameters	Essential Oils		<i>Essential Oil Association of O. basilicum</i> <i>Essential Oil (Hadipoentiyanti and Sri, 2018)</i>
	<i>O. basilicum</i> L. stem	<i>O. tenuiflorum</i> L. stem	
Organoleptic			
Shape	Liquid	Liquid	-
Colour	Brownish yellow	Pale yellow	Light yellow
Odor	Specific	Specific	-
Taste	Tighten	Bitter	-
Yield percentage	0.183% (v/w)	0.200% (v/w)	-
Refractive index	1.2906	1.3493	1.510-1.5165
Specific gravity (g/mL)	0.98054	1.0108	0.952-0.973
Solubility in absolute alcohol	1:1 (soluble)	1:1 (soluble)	4:1 (soluble)

Figure 1. Chromatogram of *O. basilicum* L. stem essential oil **Figure 2. Chromatogram of *O. tenuiflorum* L. stem essential oil**

Table 3. Chemical components of *O. basilicum* L. and *O. tenuiflorum* L. stem essential oils

Samples	Rt	Quality	Compounds	Percentage (%)
<i>O. basilicum</i> L. Stem Essential Oils	20.230	99	α -copaene	4.93
	21.559	99	β -bourbonene	1.58
	23.492	95	linalool	1.58
	25.573	99	Caryophyllene	18.75
	29.340	99	β -selinene	0.81
	30.192	97	z-citral	2.37
	31.464	99	Germacrene-D	9.19
	32.930	96	e-citral	3.39
	34.178	91	d-gurjunene	3.16
	44.959	93	Caryophyllene oxide	0.64
	47.126	98	Methyl eugenol	52.6
	49.659	98	Eugenol	2.4
	50.983	98	1,3-benzodioxole	0.75
<i>O. tenuiflorum</i> L. Stem Essential Oils	20.239	99	α -copaene	5.56
	21.568	99	β -bourbonene	1.64
	25.582	99	Caryophyllene	17.28
	29.340	99	α -Humulene (CAS)	0.76
	31.473	99	Germacrene-D	9.29
	34.187	94	Germacrene-A	3.52
	44.968	93	Caryophyllene oxide	0.86
	47.126	99	Methyl eugenol	56.72
	49.664	98	Eugenol	3.17
	50.511	97	t-Murolol	0.41
	50.983	98	1,3-benzodioxole	0.79

Note: The percentage of components is the relative percentage calculated based on the area of the peak

of linalool, eugenol, 1,8-cineol, estragol, limonene, bergamotene, α -cadinol, methyl cinnamate, and (z) cinnamic acid methyl ester (Khair-ul-Bariyah et al., 2012). In another study, it was stated that the eugenol content in the oil was quite high (10-19%), whereas methyl eugenol was a minor compound (<5%) (Zamfirache et al., 2011). All *O. basilicum* L. essential oils of the leaves from different regions in the Kingdom of Saudi Arabia contain β -Linalool (9.12-72.59%) and 1,8 Cineole (0.4-10.72%) as the main compounds. While, the content of methyl eugenol in all of the samples was recorded in the range of 0.27-18.39% (Ladwani et al., 2018). In *O. tenuiflorum* L., eugenol (25.3-51.5%) was a major constituent followed by caryophyllene (1.2-25.4%) (Sims et al., 2013). The contents of methyl eugenol as the major constituent in essential oils from whole herb, leaves, stems and inflorescences of *O. tenuiflorum* L. were 72.5%, 75.3%, 83.7% and 65.2%, respectively. Whereas, the respective concentrations of β -caryophyllene as second most dominant constituent

in the essential oils were 5.5%, 6.4%, 2.7% and 12.0% (Kothari, 2014).

Comparison of components of essential oils from stems of these two plants shows a slight similarity, both contain high amounts of methyl eugenol. Methyl eugenol is an oxygenated compound. Macchia et al. (2006) reported that methyl eugenol, produced up to 8% in the vegetative stage, was only found in some of the cultivars. Methyl eugenol is formed through cyclic acid pathways, where phenylalanine is the precursor. Through a few reaction stage, eugenol will be formed. With methyl donor from Sadenosylmethionine (SAM) and enzymatic activity from O-methyltransferase (OMT), eugenol is then converted to methyl eugenol (Murningsih et al., 2009).

The intensity of antioxidant activities of essential oils from *O. basilicum* L. and *O. tenuiflorum* L. stem was classified as very strong antioxidant (<50 μ g/mL). *Ocimum* plants contain large amounts of antioxidants,

such as flavonoids, carotenoids, vitamin C, and vitamin E. The presence of antioxidant compounds provides protection against free radical which can cause oxidative damage of cellular component (Pandey *et al.*, 2014). Natural essential oils contain several components. When they are used to protect some material, their activity is thought to be due to the dominance of the effective content of certain components therein. This is true in some cases. However, there are some exceptions. The antioxidant activity of essential oil, in fact, is also the result of the complex interplay among components and material to be protected from oxidation (Amorati, 2013).

CONCLUSION

Based on our results, there is a similarity in the chemical composition between *O. basilicum* L. and *O. tenuiflorum* L. stem essential oils. Methyl eugenol is a major compound that is produced by *Ocimum* and it might have a role in their antioxidant activities. These findings suggest that both oils might be good natural antioxidant agents.

ACKNOWLEDGEMENT

We thank to the Research Institute and Development of Universitas Muhammadiyah Prof. DR. HAMKA, Jakarta, for Funding Research Grants Development of Science and Technology Batch I 2019.

REFERENCES

- Amorati R, Foti MC & Valgimigli L, (2013), Antioxidant Activity of Essential Oils, *Journal of Agricultural and Food Chemistry*, 61(46), 10835-10847
- Baser KHC & Buchbauer G. (2010). *Handbook of essential oils: science, technology, and applications*. New York: CRC Press Taylor & Francis Group, 1-5
- Brand-Williams W, Cuvelier ME & Berset C. (1995). Use a Free Radical Method to Evaluate Antioxidant Activity. *Lebensm. Wiss. Technol.*, 28, 25-30
- Gunawan D & Mulyani S. 2004. *Ilmu Obat Alam (Farmakognosi)*. Ed I. (In Bahasa). Jakarta: Penebar Swadaya, 108-120.
- Hadipoentiyanti E & Sri W. (2008). Keragaman Selasih (*Ocimum* Spp.) Berdasarkan Karakter Morfologi, Produksi Dan Mutu Herba (In Bahasa). *Jurnal Littri*, 142-146.
- Herebian D, Choi J-H, El-Aty AMA, Shim J-H & Spiteller M. (2009). Metabolite analysis in *Curcuma domestica* using various GC-MS and LC-MS separation and detection techniques. *Biomed. Chromatogr*, 23, 951-965.
- Khair-ul-Bariyah S, Ahmed D & Ikram M. (2012). *Ocimum Basilicum*: A Review on Phytochemical and Pharmacological Studies. *Pak.J.Chem*, 2(2), 78-85.
- Kothari SK, Bhattacharya AK, Ramesh S, Garg SN, Khanuja S. (2014). Volatile Constituents in Oil from Different Plant Parts of Methyl Eugenol-Rich *Ocimum tenuiflorum* L.f. (syn. *O. sanctum* L.) Grown in South India. *Journal of Essential Oil Research*, 17(6), 656-658
- Ladwani AMA, Salman M & Abdel Hameed ES. (2018). Chemical composition of *Ocimum basilicum* L. essential oil from different regions in the Kingdom of Saudi Arabia by using Gas chromatography mass spectrometer. *Journal of Medicinal Plants Studies*, 6(1), 14-19.
- Lutony TL & Rahmayati Y. (1994). *Produksi Dan Perdagangan Minyak Atsiri*. (In Bahasa). Jakarta: Penebar Swadaya, 112-113.
- Macchia M, Pagano A, Ceccarini L, Benvenuti S, Cioni PL & Flamini G. (2006). Agronomic and phytochemical characteristics in some genotypes of *Ocimum basilicum* L. *Acta Hort.*, 723, 143-149.
- Marwat SK, Fazal-Ur-Rehman, Khan MS, Ghulam S, Anwar N, Mustafa G & Usman K. (2011). Phytochemical Constituents and Pharmacological Activities of Sweet Basil-*Ocimum basilicum* L. (*Lamiaceae*). *Asian Journal of Chemistry*, 23(9), 3773-3782.
- Mead D. (2014). *Basils (Ocimum spp.) in Indonesia*. Sulawesi: Sulang Language: Data and Working Papers; Topics in Lexicography, 3-5.
- Murningsih T, Chairul, & Kuncari ES. (2009). Methyl Eugenol, Chemotype of essential Oils of *Melaleuca* spp. (*Myrtaceae*) Growing in Cibodas Botanical Garden. (In Bahasa) *Berita Biologi*, 9(6), 809-816.
- Pandey AK, Singh P & Tripathi NN. (2014). Chemistry and Bioactivities of Essential Oils of Some *Ocimum* Species: An Overview. *Asian Pasific Journal of Tropical Biomedicine*, 4(9), 682-694.
- Ravi P, Elumalai A, Eswaraiah MC & Kasarla R. (2012). A Review on Krishna Tulsi, *Ocimum tenuiflorum* Linn. *IJRAP* 3(2), 291-293.
- Saifudin A, Rahayu V, & Teruna HY. (2011). *Standarisasi bahan obat alam*. (In Bahasa), Yogyakarta: Graha Ilmu, 90-93.

Sims CA, Juliani HR, Mentreddy SR & Simon JE. (2013). Essential Oils in Holy Basil (*Ocimum tenuiflorum* L.) as Influenced by Planting Dates and Harvest Times in North Alabama. *Journal of Medicinally Active Plants*, 2(3), 33-41.

Sulianti SB. (2008). Phytochemical Study of *Ocimum* spp.: Chemical Components of Essential Oils of Kemangi and Ruku-Ruku. (In Bahasa) *Berita Biologi*, 9(3), 237-241.

Zamfirache MM, Padurariu C, Burzo I, Olteanu Z, Boz I & Lamban C. (2011). Research Regarding the Chemical Composition of the Volatile Oil of Some Taxa Belonging to the Genus *Ocimum*, *Biologie vegetală*, 31-34.